



ORAL PIERCING

From hairstyles to clothing, tattoos to jewelry, today's youth are making a fashion statement. But those who want to look cool with oral piercing may be looking to their dentists and physicians for help afterwards.

Common symptoms after oral piercing include pain, swelling, infection, an increased flow of saliva and injuries to the gum tissue. If a blood vessel was in the path of the needle during the piercing, severe and difficult-to-control bleeding can result. For some, chipped or cracked teeth, blood poisoning or even blood clots can occur. For many, the swelling of the tongue is a common side effect. And in extreme cases, a severely swollen tongue can actually close off the airway and prevent breathing.

Unfortunately, many young people with oral piercing don't realize that these alarming side effects could happen to them. So, skip the mouth jewelry and let your healthy smile make your fashion statement!

How can oral piercing be bad for your health?

Because your mouth contains millions of bacteria, infection is a common complication of oral piercing.

Pain and swelling are other side effects of piercing. Your tongue -- a popular piercing site in the mouth -- could swell large enough to close off your airway. Piercing also can cause uncontrollable bleeding or nerve damage.

The jewelry itself also presents some hazards. You can choke on any studs, barbells or hoops that come loose in your mouth, and contact with the jewelry can chip or crack your teeth. *Source: American Dental Association.*

**For dentists and specialists in your area, use our
“Find-A-Dentist” feature at www.azda.org**

**Arizona Dental Association
3193 Drinkwater Blvd
Scottsdale AZ 85251
(480) 344-5777 or (800) 866-2732
Fax (480) 344-1442**